

Some Diophantine relations involving circular functions of rational angles

F. Calogero,

Istituto di Fisica, Università di Roma

00185 Roma, Italia

A.M. Perelomov

Institute of Theoretical and Experimental Physics

B. Cheremushkinskaya, 25

117259 Moscow, USSR

Abstract

The eigenvalues of the 3 off-diagonal matrices of rank n with elements $1 + i \cot [(j - k)\pi/n]$, $\sin^{-2}[(j - k)\pi/n]$ and $\sin^{-4}[(j - k)\pi/n]$, ($j = 1, 2, \dots, n$, $k = 1, 2, \dots, n$, $j \neq k$) are computed. The sums over k from 1 to $n - 1$ of $\cot(k\pi/n) \sin(2sk\pi/n)$ and $\sin^{-p}(k\pi/n) \cos(2sk\pi/n)$ are moreover computed for s integer and $p=2$ and 4. The results are given by simple formulae in terms of integers.

In this note we report some remarkable relations involving circular functions of rational angles.

Theorem 1. *The off-diagonal hermitian matrix of rank n whose elements are defined by the formula*

$$A_{jk} = (1 - \delta_{jk}) \left\{ 1 + i \cot \frac{(j - k)\pi}{n} \right\} \quad (1)$$

has the integer eigenvalues

$$a_s = 2s - n - 1, \quad s = 1, 2, \dots, n \quad (2)$$

and the corresponding eigenvectors $v^{(s)}$ have components

$$v_j^{(s)} = \exp\left(-\frac{2\pi i s j}{n}\right), \quad j = 1, 2, \dots, n. \quad (3)$$

Theorem 2. The 2 off-diagonal hermitian matrices whose elements are defined by the formulae

$$B_{jk} = (1 - \delta_{jk}) \sin^{-2} \frac{(j-k)\pi}{n}, \quad (4)$$

$$C_{jk} = (1 - \delta_{jk}) \sin^{-4} \frac{(j-k)\pi}{n} \quad (5)$$

are related to the matrix A defined above by the equations

$$B = \frac{1}{2} (A^2 + 2A - \sigma_n^{(1)} I), \quad (6)$$

$$C = -\frac{1}{6} (B^2 - 2(2 + \sigma_n^{(1)})B - \sigma_n^{(2)} I) \quad (7)$$

where I is the unit matrix and

$$\sigma_n^{(1)} = \frac{1}{3}(n^2 - 1), \quad \sigma_n^{(2)} = \frac{1}{45}(n^2 - 1)(n^2 + 11). \quad (8)$$

Thus their eigenvalues corresponding to the eigenvectors $v^{(s)}$ of components (3) can be written as follows:

$$b_s = \sigma_n^{(1)} - 2s(n-s), \quad s = 1, 2, \dots, n, \quad (9)$$

$$c_s = \sigma_n^{(2)} - 2s(n-s) \frac{s(n-s)+2}{3}, \quad s = 1, 2, \dots, n. \quad (10)$$

Theorem 3. The following sum rules hold:

$$\sum_{k=1}^{n-1} \cot \frac{k\pi}{n} \sin \frac{2sk\pi}{n} = n - 2s, \quad s = 1, 2, \dots, n-1, \quad (11)$$

$$\sum_{k=1}^{n-1} \sin^{-2} \frac{k\pi}{n} \cos \frac{2sk\pi}{n} = b_s, \quad s = 1, 2, \dots, n, \quad (12)$$

$$\sum_{k=1}^{n-1} \sin^{-4} \frac{k\pi}{n} \cos \frac{2sk\pi}{n} = c_s, \quad s = 1, 2, \dots, n, \quad (13)$$

$$\sum_{k=1}^{n-1} \sin^{-2p} \frac{k\pi}{n} = \sigma^{(p)} \quad (14)$$

with b_s , c_s , $\sigma_n^{(1)}$ and $\sigma_n^{(2)}$ defined above and

$$\sigma_n^{(3)} = \sigma_n^{(1)} \frac{2n^4 + 23n^2 + 191}{315}, \quad (15)$$

$$\sigma_n^{(4)} = \sigma_n^{(2)} \frac{3n^4 + 10n^2 + 227}{315}. \quad (16)$$

Theorems 1 and 2 can be easily evinced from the results of [1] or can be verified by direct computation of the eigenvalue equations by using of the explicit expression (3) for the eigenvectors. It should be emphasized that the matrix equations (6) and (7) are nontrivial consequences of the definitions (4), (5) and (1). Indeed, the validity of (6) and (7) for the diagonal elements is easily seen to correspond to the sum rules (14) with $p = 1$ and $p = 2$ using the obvious remark that

$$\sum_{k=1, k \neq j}^n \sin^{-2p} \frac{(j-k)\pi}{n} = \sum_{k=1}^{n-1} \sin^{-2p} \frac{k\pi}{n}, \quad j = 1, 2, \dots, n. \quad (17)$$

The validity of (6) for the non-diagonal terms can be verified explicitly by simple manipulations using the trigonometric identity

$$\cot \alpha \cot \beta = -1 - (\cot \alpha - \cot \beta) \cot (\alpha - \beta), \quad (18)$$

and the obvious remark

$$\sum_{k=1, k \neq j}^n \cot \frac{(j-k)\pi}{n} = \sum_{k=1}^{n-1} \cot \frac{k\pi}{n} = 0, \quad j = 1, 2, \dots, n; \quad (19)$$

while the validity of (7) for the non-diagonal terms implies the nontrivial sum rule

$$\sum_{l=1, l \neq j, k}^n \sin^{-2} \frac{(j-l)\pi}{n} \sin^{-2} \frac{(k-l)\pi}{n}$$

$$= 2(2 + \sigma_n^{(1)}) \sin^{-2} \frac{(j-k)\pi}{n} - 6 \sin^{-4} \frac{(j-k)\pi}{n}, \quad (20)$$

$$j, k = 1, 2, \dots, n, \quad j \neq k.$$

The first 3 results of Theorem 3, Eqs. (11)–(13), are obtained from the explicit computation of the eigenvalue equations, while the last result, Eq. (14), can be verified through an explicit computation of the traces of B , C , BC and C^2 from (9) and (10) and from (4), (5) using (17). Computation of the traces of AB or AC yields instead only special cases of the more general and quite trivial, formula

$$\sum_{\substack{k=1, k \neq j \\ j=1, 2, \dots, n}}^n \cot \frac{(j-k)\pi}{n} \left| \sin \frac{(j-k)\pi}{n} \right|^p = \sum_{k=1}^{n-1} \cot \frac{k\pi}{n} \left| \sin \frac{k\pi}{n} \right|^p = 0, \quad (21)$$

that holds for arbitrary (complex) p .

References

- [1] F. Calogero, A.M. Perelomov
Properties of certain matrices related to the equilibrium configuration of the one-dimensional many-body problems with the pair potentials $V_1(x) = -\log|\sin x|$ and $V_2(x) = 1/\sin^2 x$,
Commun. Math. Phys. **59**, 109–116 (1978)